

Targeted Protein Degrader

An early mover in the targeted protein degradation (TPD) space, WuXi AppTec has developed partnerships with multiple TPD players since 2016. We look forward to supporting your TPD project.

Proven Synthetic Experiences

- 158,000+ compounds synthesized
- 1,000+ experienced chemists
- 150+ clients/collaborations
- 100+ novel E3 ligase ligands achieved
- 300+ novel linkers achieved
- 300+ ADC linkers available for DAC modality
- · Custom delivery from mg to multi-kg

Leading Analytical Capabilities

- 75 dedicated separation chemists
- 200+ methods for prep-HPLC and chiral-SFC
- Diversified techniques (RP-HPLC; NP-HPLC; SFC)
- Various columns and modifiers available
- Innovative online dilution system designed for PROTACs
- Native-MS analysis for ternary complex confirmation
- Proteomics studies to identify off-target effect

Well-Established Modular System

E3 ligase ligands: Common ligands are readily available, e.g. VHL, CRBN, etc.

Linkers: Most frequently used linkers, e.g. PEG, alkyl, "clickable", rigid, etc.

PROTAC: Proteolysis Targeting Chimeras




Contact us





Research Chemistry Services

Experience with TPD Technologies



Emerging New TPD Technology: Degrader-Antibody Conjugate (DAC)

Antibody

High Specificity
 minimize off-target effect

 low immunogenicity



- Cysteine (even stoichiometry)
- Lysine (different stoichiometry)

Linker

- Cleavable
 lysosomal protease sensitive
 acid sensitive
 glutathione sensitive
- Non-cleavable

Payload/Protacs

- Highly potent small molecule toxin vs
- large and lipophilic degrader

DAC Precursor: Degrader with Cleavable Linker

